

Visualisation of radioactive particles/ diffusion cloud chamber



The goal of this experiment is to make radioactive particles visible and investigate their properties.

Physics

Modern Physics

Nuclear & particle physics

Applied Science

Medicine

Nuclear Medicine



Difficulty level

medium



Group size

2



Preparation time

45+ minutes



Execution time


45+ minutes

This content can also be found online at:



<http://localhost:1337/c/608575fd5e789700039b0f94>

PHYWE



General information

Application

PHYWE



Setup

Scientists eventually identified several distinct types of radiation, the particles resulting from radioactive decays. The three types of radiation were named after the first three letters of the Greek alphabet: (alpha), (beta), and (gamma). Alpha particles are helium nuclei, beta particles are electrons and gamma particles are photons.

This experiment tries to make these types of radioactive particles visible.

Other information (1/2)

PHYWE

Prior knowledge



The prior knowledge required can be found in the theory section.

Scientific principle



Radioactivity is a subject in our society which has been playing an important role throughout politics, economy and media for many years now. The fact that this radiation cannot be seen or felt by the human being and that the effects of this radiation are still not fully explored yet, causes emotions like no other scientific subject before.

The high-performance diffusion cloud chamber serves for making the tracks of cosmic and terrestrial radiation visible so that a wide range of natural radiation types can be identified. Furthermore, the diffusion cloud chamber offers the opportunity to carry out physical experiments with the aid of artificial radiation sources.

Other information (2/2)

PHYWE

Learning objective



The goal of this experiment is to make radioactive particles visible and investigate their properties.

Tasks



1. Determination of the amount of background radiation
2. Visualisation of α , β , γ -particles and mesons

Theory (1/8)

PHYWE

1. Natural Earth radiations

1.1 Cosmic rays

Radiations of particles with a high energy content (an exception: photon rays, which are electromagnetic waves) use to come from space down to every part of our terrestrial atmosphere (the primary cosmic radiations). Here are their main compositions:

Particles	Percentage
protons	about 90%
alpha-particles	about 9%
bigger nuclei	up to 1%

The particles penetrating into the atmosphere happen to bang into nuclei of the atmosphere and provoke nuclear reactions as well as nuclear splits. Therefore, new nuclei and elementary particles are created, go on flying and lead to other interactions.

Theory (2/8)

PHYWE

In the atmosphere layers close to earth (less than 20 km high) one can observe only one secondary kind of radiations brought up by the numerous interaction-processes in the superior layers of the atmosphere. One must differentiate four kinds of components which have a different penetrating power (see Fig. 1):

Cosmic secondary rays Components

Nucleons	protons / neutrons
Electrons and photons	electrons / positrons / photons
Mesons	mesons of different charges
Neutrinos	neutrinos / antineutrinos




Fig. 1:
Natural
splitting
processes

Theory (3/8)

PHYWE

In the Diffusion Cloud chamber, all particles charged in electricity can be detected, that is to say: electrons, positrons, mesons, protons and alpha-particles. Photons create only indirectly a trail when, by example, they eject an electron from an atom, which produces a trail of ionisation. Neutrons can lead to nuclear reaction and then, the charged particle from the nucleus creates a trail:

Particles	Symbol	Relative mass	Charge	Radioactive Period
electron	e^-	1	-1	stable
positron	e^+	1	+1	stable
myon	μ^-	206.77	-1	$1.5 \cdot 10^{-6} \text{ s}$
antimyon	μ^+	206.77	+1	$1.5 \cdot 10^{-6} \text{ s}$
proton	p^+	1836.10	+1	stable
neutron	n	1838.62	0	14.7 min
alpha-particle	α, He^{++}	7294.1	+2	stable

Theory (4/8)

PHYWE

1.2 Terrestrial radiations

All materials on Earth (earth, water, atmosphere, animals...) contain natural radionuclides which send radiations. They have been existing since the creation of Earth (that is to say 4.5 thousand million years) or are constantly being created:

U-238; Th-232; K-40 or Rb-87, for instance, belong to the natural radionuclides and have a very long radioactive period. Ra- 226; Rn-222; Po-218 or Pb-210 are constantly being created and those radionuclides have a rather short radioactive period in the three natural splitting processes. There exist also natural radionuclides with a relative short-timed radioactive period, but these do not belong to the splitting process. They are constantly being created in the upper layers of atmosphere, such as C-14 from N-14 or H-3 from N-14 or O-16, for instance.

Theory (5/8)

PHYWE

The about 100 natural radionuclides which have been existing since the creation of Earth or which are constantly being created can be found on the whole Earth in different concentrations. That is the reason why there are constant exchanges between earth, water, atmosphere and animal-life (Fig. 2).

That is why the Diffusion Cloud chamber is given materials made of natural radionuclides, which emit radiations.




Fig. 2

Theory (6/8)

PHYWE

2. The Diffusion Cloud chamber

How are tracks formed in the Cloud chamber?

The particles of which the radiation consists are unimaginably small. As an example, if we think of 1 trillion (1,000,000,000,000) protons lined up one after the other, they would only give a 2 mm long line! The particles in radiation can therefore not even be seen under the best microscopes.

Under certain conditions, however, the particles can produce cloud tracks which are visible to the human eye. Such tracks enable us to see where the particles have travelled. This is similar to looking at an aeroplane which is flying at a high altitude. The aeroplane itself is not visible from the ground, only the condensation trail can be seen.

Theory (7/8)

PHYWE

Alpha particles, electrons, protons and mesons (each of the particles named carries an electrical charge) all produce cloud tracks in a cloud chamber. (The tracks of the individual particles look different, so that one can determine,

- which particle has flown through the cloud chamber,
- the speed at which it has flown (energy) and
- whether it had a collision or was deflected during its flight.

The alcohol vapour diffusing down from above to the black plate liquefies (condenses to drops), as soon as it reaches the immediate vicinity of the cooled plate.

Theory (8/8)

PHYWE




Fig. 3: The creation of cloud tracks in a cloud chamber.

Above the liquefied alcohol vapour, there is a 1 - 2 mm thick layer in which the vapour has just not quite liquefied. In this layer, drop formation, and so cloud formation, can be deliberately caused, e.g. by finest dust particles (condensation nuclei) or by a radiation particle flying through. During their flight, radiation particles "damage" (ionize) numerous alcohol molecules, which can then take on very much bigger alcohol drops and so become visible to us. They form the cloud track (Fig. 3).

Equipment

Position	Material	Item No.	Quantity
1	Diffusion cloud chamber, 45 x 45 cm PJ45, 230 VAC	09046-93	1
2	Isopropyl alcohol, extra pure, 1000 ml	30092-70	2

PHYWE



Setup and Procedure

Setup and Procedure (1/6)

PHYWE

The cloud chamber consists of a chamber base and the observation chamber. The chamber base comprises a cooling element, a power supply, an alcohol reservoir, an alcohol pump and a programmable time switch. The observation chamber is placed onto the chamber base.

The bottom of the observation chamber is formed by a massive, black metal plate (surface 45 cm×45 cm) which is cooled over the whole surface to about -30°C by means of the cooling element.



Fig. 4: Experimental setup

Setup and Procedure (2/6)

PHYWE

The top plate and the side plates of the observation chamber consist of two glass hoods which are placed one into the other. A grid of fine heating wires is placed between the upper two glass plates. These wires serve for heating this area of the chamber thus keeping the hood free from condensation. At the same time, they are held at high voltage thus generating an electric field that attracts the ions. The upper part of the glass hood includes an electrically heated gutter which runs around the whole circumference. Iso-propyl alcohol flows through a bended tube and drops into the gutter.

The alcohol evaporates and diffuses from the upper, warmer area of the chamber to the cold chamber bottom. There the alcohol is condensed into tiny droplets and flows back into the reservoir.

Right above the thin liquid layer covering the bottom a zone of oversaturated alcohol vapour is formed. It's in this area, and only in this area, that the charged material particles coming from the inside or from the outside produce ions along their trajectory. The tiny alcohol droplets preferably attach to these ions thus producing a visible cloud track. The length and the structure of the cloud track give information on the kind of ionizing particles.

Setup and Procedure (3/6)

PHYWE

In order to guarantee an optimum view on the observation chamber, it is recommended to place the chamber onto a square table (border length about 90 to 100 cm) which should be about 30-60 cm high.

Please make sure that the ventilating slots are not covered and that the cloud chamber is not subjected to direct light coming from above. A slightly darkened room would be perfect. Use the connecting cable supplied to connect the cloud chamber to mains. The mains connection is located down on the back side. The corresponding mains socket should be protected for 10 A to 16 A.

Adjust the cloud chamber on an absolutely horizontal level with the aid of the adjustable feet in order to provide an even alcohol level in the gutter and thus a clear image.

When you have connected the cloud chamber to mains and opened the front side, remove the alcohol reservoir from its installation place and unscrew the union nut of the double tubing assembly. Refill the alcohol reservoir, fix the double tubing again and place the reservoir back in place.

Setup and Procedure (4/6)

PHYWE

Now activate the switches as follows:

main switch	ON
mode	continuous operation
high voltage	ON

The knurled nut serves for regulating the amount of alcohol dropping into the evaporating gutter. Turn the knurled nut to the left and observe the alcohol flowing through the bended tube and dropping into the gutter. When the gutter is filled to a liquid level of about 1 cm, reduce the alcohol supply to about 2 droplets per second. When the cloud chamber is being operating, the alcohol in the gutter should remain constant at this level. After about 5 minutes, the first white tracks should appear on the black observation surface. If, however, after 1 hour the tracks become a bit fuzzy and milky, you have to reduce the gutter heating by means of the control knob.

Setup and Procedure (5/6)

PHYWE

If the tracks are too weak, the heating of the gutter must be increased.

If you want to run the cloud chamber at automatic operation, set the markers of the programmable time switch to the desired switching time (red marker to switch the cloud chamber on, green marker to switch it off) and set the mode switch to "timer". For more detailed information see the instructions on the programmable time switch.

Now close the front side by inserting the plate into the right side of the opening, pushing it back to the left stop position and locking it up.




Fig. 5: Section through diffusion cloud chamber.

Setup and Procedure (6/6)


PHYWE

Artificial radiation sources

On the left side of the chamber base there is an opening which serves for introducing artificial radiation sources. Use the head of the screw to push the plate to the right. The ball with pin which is located behind the plate can be turned around with the aid of the pin until the opening towards the inner area becomes visible. Now you can insert an artificial radiation source like, for instance, the thorium source a few centimeters into the inner area of the chamber.

PHYWE

Evaluation



Task 1 (1/2)

PHYWE

All of the radioactive particles described in the previous chapter are continually present as so-called background radiation, unless one is in a special screened off room. When radioactive effects such as, for example, radioactive decay, are studied, then the zero rate resulting from the background radiation which was prevailing at that time must always be subtracted from the effects under observation. In general, the zero rate is about 18 registered impulses per minute. This measurement is usually made by means of a tube counter which is connected to a counting instrument. A glance at the active area of the cloud chamber shows, however, an apparently very great abundance of particles, which one tends to associate rather with the presence of a radioactive material in the chamber than merely to the background radiation. A little trick helps here to categorize the density of the particle tracks as really being due to background radiation.

Task 1 (2/2)

PHYWE

Take a piece of typing paper and cut out a hole of about 0.8 cm diameter in the middle of it. Lay the piece of paper flat on the glass plate of the cloud chamber. Now, from a distance of about 10 cm from the paper, look through the hole with one eye and observe the active area of the cloud chamber. Count twenty "parts" of particle tracks which are visible through the hole and measure the time elapsed while doing so. This trick with the hole in the paper simulates the opening of a tube counter. Only those particles which hit the tube counter opening can be registered. From the multitude of particle tracks in the cloud chamber, only those are now visible which pass through an opening which corresponds to that of a tube counter. The zero rate which is theoretically to be expected can be so confirmed.

Task 2 (1/8)

PHYWE

In the chamber, one can observe tracks of “clouds” which are generated by α -particles, protons, electrons/positrons and mesons.

a) Example 1

Most of time, one can observe short tracks and thinner, longer tracks. To begin with, let us concentrate on short, but the biggest tracks (Fig. 6).

The appearance of α -tracks on the whole observation place is statistically dispersed, that is the reason why one cannot foretell when and where the next track will appear.



Fig. 6: Track generated by an α -particle.

Task 2 (2/8)

PHYWE

In the air, α -particles measure about 5 cm but in the steam of alcohol they are not longer. Moreover, α -particles can be absorbed by a single sheet of paper. How can they reach the interior of the Cloud chamber?

On the one hand, α -particles may have been liberated in the chamber itself by a radioactive nucleus. On the other hand, it may be protons with a high content of energy which were formed during the secondary radiations process in the atmosphere. They can penetrate into the chamber across the glas protection. After penetrating the chamber, if their energy content is low enough as to being able to give their energy to the atom-electrons of the gas inside the chamber, they generate a track, which looks like the one of the α -particles (Fig. 7).




Fig. 7: Track generated by protons.

Task 2 (3/8)

PHYWE

When the particle penetrate vertically the layer of supersaturated Alcohol steam, one can recognize only a spot as an ionisation trace.

b) Example 2

When the observer now observes the thin and to some extend very longer tracks (tracks with low drops density), one will be puzzled by the large amount of track manifestations. Therefore, we recommend you to consider precise forms – that is to say: the length of those tracks.

- First of all, observers should try to recognize a thin, straight and long trajectory – across the whole observation place. This shows particularly fast (with a high content of energy) electrons (Fig. 8).
- Slowly flowing electrons (i.e. with a low content of energy) have shorter trajectories, which are partly curved or buckled (Fig. 9) because of deviation.

Task 2 (4/8)

PHYWE

- Electrons with a very low content of energy generate short trajectories, which look ornated or tortuous because of many deviations of the atoms in the steam layer (Fig. 10).
- Provided you place a powerful Ra-emitter, the γ -quanta will penetrate the chamber across the glass and release a large amount of photon-electrons or compton-electrons. Then, a formation of short and many times tortuous “worm-like” trajectories appears on the whole surface (Fig. 11).




Fig. 8: Track generated by an electron with a high content of energy.

Task 2 (5/8)

PHYWE




Fig. 9: Track generated by repeated deviated β -particle.



Fig. 10: Tracks of β -particles with a low content of energy.

Task 2 (6/8)

PHYWE




Fig. 11: Track generated by protons-electrons and compton electrons.

c) Example 3

Mesons, which represent 90% of the secondary cosmic rays can be detected in the Cloud chamber too. m-mesons take an important part in this process since they have a positive or a negative elementary charge and their weight corresponds to 207 times electrons weight.

Mesons with a high content of energy create trajectories which look like tracks generated by electrons. However, high moderated mesons ionize highly and produce tracks which correspond nearly to those of α -particles. Therefore, it will be very difficult, in individual cases, to know whether one is being observing α -particles, protons, mesons or electrons (Fig. 12).

Task 2 (7/8)

PHYWE

Since the average lifetime of μ -mesons corresponds to micro-seconds, one can occasionally observe the dissociation of a μ -meson into an electron and into two invisible neutrinos in the Cloud chamber. The electron does not move to the same direction as the meson before, so that one can observe a characteristic bend in the track (Fig. 13). However, after the collision, one can clearly observe a thinner track, since the electron has a smaller ionisation density than the meson. The number of generated pairs of ions in each trajectory (in comparable conditions) are as follows:

- α -particles: 10.000
- protons: 5.000
- electrons: 100
- gamma-quanta: 1 (specific ionisation).

Task 2 (8/8)

PHYWE




Fig. 12: Track generated by a meson.




Fig. 13: Track of dissociation of a μ -meson into an electron and two neutrinos.